arXiv:0710.0563v1 [hep-ph] 2 Oct 2007

Newly observed charmed states: the case of $X(3872)$

Stefano Nicotri

Università degli studi di Bari and INFN sezione di Bari,
via Orabona 4, I-70126, Bari, Italy
`stefano.nicotri@ba.infn.it`

Summary. I briefly review the properties of the hidden charm meson $X(3872)$, discussing some puzzling aspects concerning its theoretical interpretation.

Introduction

Recently many new mesons have been observed in the open and hidden charm sectors [1, 2]. From the need of theoretical interpretation of those states, the interesting possibility of identifying new "exotic" structures has been pointed out. In the following I discuss the case of $X(3872)$, which represents one of the most interesting examples in this sense. In particular I focus the attention on the properties of the radiative decays $X \to D\bar{D}\gamma$ [3].

$X(3872)$

The experimental observations can be summarized as follows:

- the $X(3872)$ resonance has been found in $J/\psi\pi^+\pi^-$ distribution by four experiments, both in B decays($B^{-(0)} \to K^{-(0)}X$), both in $p\bar{p}$ annihilation [4]. The mass is $M = 3871.9 \pm 0.6$ MeV while the width remains unresolved: $\Gamma < 2.3$ MeV (90 % CL);
- there is no evidence of resonances in the charged mode $J/\psi\pi^\pm\pi^0$ or in $J/\psi\eta$ [5];
- the state is not observed in e^+e^- annihilation;
- for X produced in B decays the ratio $\frac{B(B^0\to K^0X)}{B(B^+\to K^+X)} = 0.61 \pm 0.36 \pm 0.06$ is obtained [5];
- the dipion spectrum in $J/\psi\pi^+\pi^-$ is peaked at large mass [6];
- the decay in $J/\psi\pi^+\pi^-\pi^0$ is observed [7] with $\frac{B(X\to J/\psi\pi^+\pi^-\pi^0)}{B(X\to J/\psi\pi^+\pi^-)} = 1.0 \pm 0.4 \pm 0.3$: this implies G-parity violation;

- the radiative mode $X \to J/\psi\gamma$ is found [7, 8] with $\frac{B(X\to J/\psi\gamma)}{B(X\to J/\psi\pi^+\pi^-)} = 0.19 \pm 0.07$, therefore charge conjugation of the state is C=+1;
- the angular distribution of the final state is compatible with the spin-parity assignment $J^P = 1^+$ [9];
- there is a signal in $D^0\bar{D}^0\pi^0$ with $\frac{B(X\to D^0\bar{D}^0\pi^0)}{B(X\to J/\psi\pi^+\pi^-)} = 9 \pm 4$ [10]; the near-threshold $D^0\bar{D}^0\pi^0$ enhancement in $B \to D^0\bar{D}^0\pi^0 K$has a the peak at $M = 3875.4 \pm 0.7^{+1.2}_{-2.0}$ MeV with $B(B \to KX \to KD^0\bar{D}^0\pi^0) = (1.27 \pm 0.31^{+0.22}_{-0.39}) \times 10^{-4}$ [11] .

All the measurements are compatible with the assignment $J^{PC} = 1^{++}$, but if the 2^3P_1 is identified with $Y(3940)$, another recently observed meson, there is an overpopulation of 1^{++} $c\bar{c}$ mesons. This, together with the observation of the G-parity violating X transitions suggested the conjecture that $X(3872)$ is not a charmonium $\bar{c}c$ state. Indeed, the coincidence between the X mass as averaged by PDG and the $D^{*0}\overline{D}^0$ mass inspired the proposal that $X(3872)$ could be a molecular quarkonium [12], a D^{*0} and $\overline{D}^0$ bound state with small binding energy due to a single pion exchange [13]. In this scheme the wave function of $X(3872)$ is given by [14]:

$$|X(3872)>= a\,|D^{*0}\bar{D}^0 + \bar{D}^{*0}D^0> +b\,|D^{*+}D^- + D^{*-}D^+> + \ldots \qquad (1)$$

(with $|b| \ll |a|$) and so one could explain why this state seems not to have definite isospin, why the mode $X \to J/\psi\pi^0\pi^0$ was not found, and why, if the binding mechanism is a single pion exchange (which is in fact a controversial point), there are no $D\overline{D}$ molecular states. The opposing remark concerning the large value of the ratio $\frac{B(X\to J/\psi\pi^+\pi^-\pi^0)}{B(X\to J/\psi\pi^+\pi^-)}$ is that the ratio of the amplitudes is small: $\frac{A(X\to J/\psi\rho^0)}{A(X\to J/\psi\omega)} \simeq 0.2$, so that the isospin violating amplitude is 20% of the isospin conserving one, an effect that could be related to the mass difference between neutral and charged D mesons, considering the contribution of DD^* intermediate states to X decays [15, 16]. Moreover, the central value of the mass measured in $D^0\bar{D}^0\pi^0$ is 4 MeV higher than the PDG value (with a large systematic error), a result difficult to explain in the molecular picture [17]. It has also been suggested that the molecular interpretation would imply that the radiative decay in neutral D mesons: $X \to D^0\bar{D}^0\gamma$ should be dominant with respect to $X \to D^+D^-\gamma$ [14]. However, assuming that $X(3872)$ is an ordinary $J^{PC} = 1^{++}$ charmonium and describing the $X(3872) \to D\bar{D}\gamma$ amplitude by polar diagrams with D^* and $\psi(3770)$ as intermediate particles the ratio $R = \frac{\Gamma(X\to D^+D^-\gamma)}{\Gamma(X\to D^0\overline{D}^0\gamma)}$ depends only on the ratio $c/\hat{g}_1$, c being the parameter governing the radiative $X\psi(3770)\gamma$ matrix element, and $\hat{g}_1$ representing the coupling $XD\bar{D}^*$ (or $XD^*\bar{D}$). The ratio R is plotted in fig.1 versus $c/\hat{g}_1$ and it is shown that in any case $R < 0.7$. Thus, there is a suppression of the radiative X decay mode into charged D mesons with respect to the mode with neutral D. Moreover, for small values of $c/\hat{g}_1$ the ratio R is tiny, so that this seems not a peculiar feature of a molecular structure of $X(3872)$.



Fig. 1. Ratio of charged $X \to D^+D^-\gamma$ to neutral $X \to D^0\bar{D}^0\gamma$ decay widths versus the ratio of hadronic parameters $c/\hat{g}_1$.



Fig. 2. Photon spectrum (in arbitrary units) in $X \to D^0\bar{D}^0\gamma$ (top) and $X \to D^+D^-\gamma$ (bottom) decays for values of the hadronic parameter $c/\hat{g}_1 = 1$ (left) and $c/\hat{g}_1 = 300$ (right).

The photon spectrum in radiative X decays to both neutral and charged D meson pairs for two representative values of $c/\hat{g}_1$, namely $c/\hat{g}_1 = 1$ and $c/\hat{g}_1 = 300$, is depicted in fig.2. For small values of the parameter $c/\hat{g}_1$, i.e. in the condition where the intermediate D^* dominates the decay amplitude, the photon spectrum in the $D^0\bar{D}^0\gamma$ mode essentially coincides with the line corresponding to the D^* decay at $E_\gamma \simeq 139$ MeV and width determined by the D^* width. The narrow peak is different from the line shape expected in a molecular description, which is related to the wave function of the two heavy mesons bound in the $X(3872)$, in particular to the binding energy of the system, being broader for larger binding energy. On the other hand, the photon spectrum in the charged $D^+D^-\gamma$ mode is broader, with a peak at $E_\gamma \simeq 125$ MeV, the total $X \to D^+D^-\gamma$ rate being severely suppressed with respect to the $X \to D^0\bar{D}^0\gamma$ one.

At the opposite side of the $c/\hat{g}_1$ range, where $\psi(3770)$ gives a large contribution to the radiative amplitude, a peak at $E_\gamma \simeq 100$ MeV appears both in neutral and charged D meson modes, in the first case together with the structure at $E_\gamma \simeq 139$ MeV. This spectrum was described also in [14], where in this case the radiative decay was interpreted as deriving from the $\bar{c}c$ core of $X(3872)$. In this range of parameters the ratio of the $X \to D^+D^-\gamma$ to $X \to D^0\bar{D}^0\gamma$ rates reaches the largest value.

The experimental determination of the photon spectrum of the type depicted in fig.2, together with the measurement of the $X \to D\bar{D}\gamma$ widths is a challenging task. Nevertheless, this measurement is important to shed light on the structure of $X(3872)$. Other methods to distinguish between a molecular and a quarkonium description have been proposed, and concern the pionic X decays to χ_{cJ} [18].

Conclusions

I have summarized the features of $X(3872)$, discussing in particular possible tests of the molecular interpretation of this state. I have focused the attention on the radiative decays $X \to D\bar{D}\gamma$, showing that the suppression of the charged channel with respect to the neutral one is not a distinctive feature of the molecular picture. However, in this decay mode the photon spectrum is expected to be different in the $c\bar{c}$ and molecular description, so that its experimental investigation could be useful to shed more light on this puzzling hadron.

Acknowledgements

I thank the conference organizers and convenors. I also acknowledge P. Colangelo and F. De Fazio for collaboration. This work has been supported in part by the EU Contract No. MRTN-CT-2006-035482, "FLAVIAnet".

References

[1] W. M. Yao *et al.* [Particle Data Group], J. Phys. G **33** (2006) 1.

[2] For reviews see:P. Colangelo *et al.* Mod. Phys. Lett. A **19** (2004) 2083; E. S. Swanson, Phys. Rept. **429** (2006) 243. P. Colangelo *et al.* arXiv:hep-ph/0609240; J. L. Rosner, J. Phys. Conf. Ser. **69** (2007) 012002 and P. Colangelo *et al.* arXiv:0706.0053 [hep-ph].

[3] P. Colangelo, F. De Fazio and S. Nicotri, Phys. Lett. B **650** (2007) 166.

[4] S. K. Choi *et al.* [Belle Collaboration], Phys. Rev. Lett. **91** (2003) 262001; D. Acosta *et al.* [CDF II Collaboration], Phys. Rev. Lett. **93** (2004) 072001; V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **93** (2004) 162002; B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **71** (2005) 071103.

[5] D. Bernard, talk at the International Conference on QCD and Hadronic Physics, June 16-20 2005, Beijing, China.

[6] A. Abulencia *et al.* [CDF Collaboration], Phys. Rev. Lett. **96** (2006) 102002.

[7] K. Abe *et al.* [Belle Collaboration], arXiv:hep-ex/0505037.

[8] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **74** (2006) 071101

[9] K. Abe *et al.* [Belle Collaboration], arXiv:hep-ex/0505038; A. Abulencia *et al.* [CDF Collaboration]. arXiv:hep-ex/0612053.

[10] G. Gokhroo et al., [Belle Collaboration], BELLE-CONF-0568 (2005); P. Pakhlov, talk at ICHEP 2006, 26/7-2/8 2006, Moskow, Russia.

[11] G. Gokhroo *et al.*, Phys. Rev. Lett. **97** (2006) 162002, The signal has been confirmed by BaBar Collaboration, see P. Grenier, arXiv: 0705.2432.

[12] M. B. Voloshin and L. B. Okun, JETP Lett. **23** (1976) 333.

[13] F. E. Close and P. R. Page, Phys. Lett. B **578** (2004) 119; M. B. Voloshin, Phys. Lett. B **579** (2004) 316; N. A. Tornqvist, Phys. Lett. B **590** (2004) 209; C. Y. Wong, Phys. Rev. C **69** (2004) 055202; E. Braaten and M. Kusunoki, Phys. Rev. D **69** (2004) 074005; E. S. Swanson, Phys. Lett. B **588** (2004) 189 and **598** (2004) 197; M. T. AlFiky *et al.* Phys. Lett. B **640** (2006) 238; E. Braaten *et al.* arXiv:hep-ph/0702128; S. Fleming*et al.* Phys. Rev. D **76** (2007) 034006.

[14] M. B. Voloshin, Int. J. Mod. Phys. A **21** (2006) 1239.

[15] M. Suzuki, Phys. Rev. D **72** (2005) 114013.

[16] C. Meng and K. T. Chao, Phys. Rev. D **75** (2007) 114002.

[17] An argument in favour of the possible coexistence of a $D^0\bar{D}^0\pi^0$ signal with the other features of the molecular description has been provided by C. Hanhart *et al.* Phys. Rev. D **76** (2007) 034007.

[18] S. Dubynskiy and M. B. Voloshin, arXiv:0709.4474 [hep-ph].